

Ancient Duplications and Expression Divergence in the Globin Gene Superfamily of Vertebrates: Insights from the Elephant Shark Genome and Transcriptome

Juan C. Opazo,^{*,†,1} Alison P. Lee,^{†,2} Federico G. Hoffmann,^{†,3,4} Jessica Toloza-Villalobos,¹ Thorsten Burmester,⁵ Byrappa Venkatesh,^{*,2,6} and Jay F. Storz^{7,*}

¹Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile

²Institute of Molecular and Cell Biology, Comparative and Medical Genomics Laboratory, Agency for Science, Technology, and Research Biopolis, Singapore

³Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University

⁴Institute for Genomics, Biocomputing, and Biotechnology, Mississippi State University

⁵Institute of Zoology, University of Hamburg, Hamburg, Germany

⁶Departments of Paediatrics, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

⁷School of Biological Sciences, University of Nebraska, Lincoln

[†]These authors contributed equally to this work.

*Corresponding author: E-mail: jopazo@gmail.com; mcbbv@imcb.a-star.edu.sg; jstorz2@unl.edu.

Associate editor: Naruya Saitou

Abstract

Comparative analyses of vertebrate genomes continue to uncover a surprising diversity of genes in the globin gene superfamily, some of which have very restricted phyletic distributions despite their antiquity. Genomic analysis of the globin gene repertoire of cartilaginous fish (Chondrichthyes) should be especially informative about the duplicative origins and ancestral functions of vertebrate globins, as divergence between Chondrichthyes and bony vertebrates represents the most basal split within the jawed vertebrates. Here, we report a comparative genomic analysis of the vertebrate globin gene family that includes the complete globin gene repertoire of the elephant shark (*Callorhynchus milii*). Using genomic sequence data from representatives of all major vertebrate classes, integrated analyses of conserved synteny and phylogenetic relationships revealed that the last common ancestor of vertebrates possessed a repertoire of at least seven globin genes: single copies of *androglobin* and *neuroglobin*, four paralogous copies of *globin X*, and the single-copy progenitor of the entire set of vertebrate-specific globins. Combined with expression data, the genomic inventory of elephant shark globins yielded four especially surprising findings: 1) there is no trace of the *neuroglobin* gene (a highly conserved gene that is present in all other jawed vertebrates that have been examined to date), 2) *myoglobin* is highly expressed in heart, but not in skeletal muscle (reflecting a possible ancestral condition in vertebrates with single-circuit circulatory systems), 3) elephant shark possesses two highly divergent *globin X* paralogs, one of which is preferentially expressed in gonads, and 4) elephant shark possesses two structurally distinct α -*globin* paralogs, one of which is preferentially expressed in the brain. Expression profiles of elephant shark globin genes reveal distinct specializations of function relative to orthologs in bony vertebrates and suggest hypotheses about ancestral functions of vertebrate globins.

Key words: Chondrichthyes, gene duplication, gene family evolution, globin, myoglobin, neuroglobin.

Introduction

Analyses of vertebrate genomes have led to the discovery of many new members of the globin gene superfamily (Hankeln et al. 2005; Hankeln and Burmester 2008; Burmester and Hankeln 2009, 2014; Storz et al. 2011, 2013). The canonical globin gene repertoire of jawed vertebrates (gnathostomes) comprises eight main members, each of which may be represented by multiple, structurally distinct paralogs with distinct expression domains that are specific to different tissues or cell types: neuroglobin (*Ngb*), cytoglobin (*Cygb*), androglobin (*Adgb*), globin E (*GbE*), globin X (*GbX*), globin Y (*GbY*), myoglobin (*Mb*), and the α - and β -type subunits of hemoglobin (*Hb*). All bony vertebrates (Euteleostomi, which

includes osteichthyans + tetrapods) that have been examined to date possess copies of *Ngb*, *Cygb*, and *Adgb* (Burmester et al. 2000, 2002, 2004; Awenius et al. 2001; Pesce et al. 2002; Trent and Hargrove 2002; Fuchs et al. 2004, 2005; Kugelstadt et al. 2004; Wystub et al. 2004; Hankeln et al. 2005; Roesner et al. 2005; Hankeln and Burmester 2008; Burmester and Hankeln 2009, 2014; Hoffmann, Opazo, et al. 2010; Kakar et al. 2010; Hoogewijs et al. 2012; Schwarze and Burmester 2013). In bony vertebrates, the monomeric *Ngb* protein is expressed in the neurons of the central and peripheral nervous system, and in some endocrine tissues, whereas the homodimeric *Cygb* protein is expressed in fibroblasts and related cell types and in distinct nerve cells in the central

and peripheral nervous systems (reviewed by Hankeln and Burmester 2008; Burmester and Hankeln 2009, 2014). The more recently discovered *Adgb* gene is especially enigmatic, as it is a chimeric fusion gene; the encoded protein has an N-terminal calpain-like domain, an internal globin domain that has undergone an internal shuffling of α -helical subdomains, and an IQ calmodulin-binding motif (Hoogewijs et al. 2012). In mammals, the *Adgb* gene is preferentially expressed in testis (Hoogewijs et al. 2012). The heme coordination chemistries and other structural features of *Ngb*, *Cygb*, and *Adgb* suggest that these globins may perform redox-regulated signaling functions or oxygen-sensing functions that mediate oxygen-dependent protein activities, but the primary physiological functions of these globin proteins are still mostly shrouded in mystery (Burmester and Hankeln 2014).

In the majority of bony vertebrate taxa, tetrameric Hb and monomeric Mb play key roles in the maintenance of cellular oxygen supply to fuel aerobic metabolism. The proto-*Hb* gene and the single-copy ancestor of the *Mb/GbE* gene pair were paralogous products of a whole-genome duplication event that occurred in the stem lineage of vertebrates (Hoffmann, Opazo, Storz, et al. 2012). A subsequent tandem duplication of the proto-*Hb* gene gave rise to the progenitors of the α - and β -globin gene subfamilies. This tandem gene duplication event occurred in the stem lineage of gnathostomes prior to the split between cartilaginous fish and bony vertebrates (Hoffmann, Opazo, Storz, et al. 2012) which is estimated to have occurred approximately 450–500 Ma in the early Paleozoic (Hedges 2009; Inoue et al. 2010). The $\alpha_2\beta_2$ Hb tetramer is most highly expressed in erythrocytes and is responsible for transporting oxygen in arterial blood from the respiratory surfaces (lungs, gills, or skin surface) to the cells of respiring tissues, and for transporting carbon dioxide in the venous blood from the tissues back to the gas exchange surfaces. In addition to the familiar respiratory function of Hb in blood-gas transport, recent reports of Hb expression in nonerythroid cell types suggest possible “moonlighting” functions that may involve the scavenging of reactive oxygen species (Nishi et al. 2008) or regulation of nitric oxide signaling (Straub et al. 2012). Mb is primarily expressed in myocytes of cardiac and skeletal muscle, where it regulates cellular oxygen tension and the bioavailability of nitric oxide (Wittenberg and Wittenberg 2003; Helbo et al. 2013). Similar to the case with Hb, recent discoveries of Mb expression in vascular smooth muscle and endothelial cells suggest that this protein may play a more versatile role than previously thought (Qiu et al. 1998; Wittenberg and Wittenberg 2003; Cossins et al. 2009; Helbo et al. 2013).

In addition to the *Ngb*, *Cygb*, *Adgb*, *Hb*, and *Mb* genes that have been retained in all or nearly all bony vertebrate lineages, a number of paralogous globins have been discovered that have far more restricted phyletic distributions. For example, one or more copies of *GbX* have been documented in lampreys and most examined bony vertebrates other than mammals and archosaurs (Roesner et al. 2005; Fuchs et al. 2006; Dröge and Makalowski 2011; Blank and Burmester 2012; Schwarze and Burmester 2013; Schwarze et al. 2014). Bony vertebrate *GbX* is a monomeric, membrane-bound globin

that possesses N-terminal acylation sites, which suggests possible roles in cellular signaling and/or protection against the oxidation of membrane lipids (Blank, Wollberg, et al. 2011; Blank and Burmester 2012). The *GbY* gene has been documented in the genomes of lobe-finned fishes (coelacanth, *Latimeria chalumnae*) and several tetrapod taxa such as the anole lizard (*Anolis carolinensis*) and platypus (*Ornithorhynchus anatinus*), but the gene does not appear to be present in lampreys, teleost fishes, birds, or therian mammals (marsupials and eutherians) (Hoffmann, Opazo, et al. 2010; Hoffmann, Storz, et al. 2010; Hoffmann et al. 2011; Schwarze and Burmester 2013; Schwarze et al. 2014). In adult *Xenopus*, *GbY* is expressed in a broad range of tissue types (Fuchs et al. 2006), but its physiological function has yet to be elucidated. The *GbE* gene was initially found only in birds (Kugelstadt et al. 2004; Hoffmann, Opazo, et al. 2010; Blank, Kiger, et al. 2011; Hoffmann et al. 2011). The monomeric *GbE* protein appears to perform a Mb-like function in regulating oxygen supply to photoreceptor cells in the avascular avian retina (Blank, Kiger, et al. 2011), although a role in regulating cellular redox homeostasis is also possible. Although initial genomic surveys suggested that *GbE* represented a bird-specific globin, phylogenetic analyses combined with assessments of conserved synteny revealed that *GbE* and *Mb* represent the paralogous products of a tandem gene duplication that occurred in the stem lineage of gnathostomes (Hoffmann et al. 2011; Hoffmann, Opazo, Storz, et al. 2012). The antiquity of the gene suggested the possibility that orthologs of *GbE* might be found in nonavian vertebrate taxa that had yet to be surveyed. Sure enough, a recent comparative genomic study identified orthologs of the hitherto “bird-specific” globin in the coelacanth (Schwarze and Burmester 2013).

The *Ngb*, *Adgb*, and *GbX* genes originated prior to the split between protostomes and deuterostomes, and were inherited by both descendant lineages (Burmester et al. 2000; Blank and Burmester 2012; Hoffmann, Opazo, Hoogewijs, et al. 2012; Hoogewijs et al. 2012; Burmester and Hankeln 2014). Vertebrate *Ngb* shows phylogenetic affinities with a diversity of nerve globins in annelid worms, echinoderms, and cephalochordates (amphioxus), and vertebrate *GbX* shows affinities with an equally diverse set of putatively N-acylated globins in protostomes and nonvertebrate deuterostomes (Blank and Burmester 2012; Hoffmann, Opazo, Hoogewijs, et al. 2012). The remaining members of the vertebrate globin gene repertoire are products of vertebrate-specific gene duplications or whole-genome duplications (Hoffmann, Storz, et al. 2010; Hoffmann et al. 2011; Storz et al. 2011, 2013; Hoffmann, Opazo, Hoogewijs, et al. 2012; Hoffmann, Opazo, Storz, et al. 2012). Cyclostomes (jawless fishes represented by lampreys and hagfish), the sister group of gnathostomes, possess a surprisingly diverse repertoire of globin genes, but available data indicate that *Cygb* is the only vertebrate-specific globin shared between cyclostomes and gnathostomes (Schwarze et al. 2014). Analyses of available globin sequences from sharks and rays have demonstrated that cartilaginous fish and bony vertebrates share orthologous α - and β -type *Hb* genes (Hoffmann, Opazo,

Hoogewijs, et al. 2012; Hoffmann, Opazo, Storz, et al. 2012; Opazo et al. 2013), but it remains possible that cartilaginous fish and cyclostomes share orthologs of other globin types that were secondarily lost in the ancestor of bony vertebrates.

It is possible that additional globin genes still await discovery as full genome sequences become available for more vertebrate taxa. Among vertebrates, it should be especially enlightening to characterize the complete globin gene repertoire for a representative of Chondrichthyes (cartilaginous fish), as divergence between Chondrichthyes and bony vertebrates represents the most basal split within the gnathostomes. Comparative expression analyses of orthologous genes shared between cartilaginous fish and bony vertebrates could potentially shed light on ancestral functions and could reveal whether certain physiological functions of modern-day globins represent derived, “repurposed” modifications of ancestral functions. Here, we report a comparative genomic analysis that includes the globin gene repertoire of the elephant shark (*Callorhinchus milii*), the first cartilaginous fish to have its full genome sequenced (Venkatesh et al. 2007, 2014). The main objectives of the study were 1) to unravel the duplicative history of globin genes during the early evolution of vertebrates, 2) to reconstruct the globin gene repertoire of the vertebrate common ancestor, and 3) to test whether orthologous globins shared between cartilaginous fish and bony vertebrates have conserved patterns of tissue-specific expression.

Results and Discussion

Using gene and genome sequences from representatives of all major vertebrate classes (Materials and Methods), we integrated analyses of conserved synteny and phylogenetic relationships to unravel the evolutionary history of vertebrate globin genes. Combined with expression data, the inventory of globin genes in the elephant shark genome yielded four especially surprising findings: 1) there is no trace of the *Ngb* gene (a highly conserved gene that is possessed by all other gnathostome taxa that have been examined to date), 2) *Mb* is highly expressed in heart, but not in skeletal muscle, 3) elephant shark possesses two highly divergent *GbX* paralogs, one of which is preferentially expressed in gonads (ovaries and testis), and 4) elephant shark possesses two structurally distinct α -globin paralogs, one of which is preferentially expressed in the brain. The divergent expression domains of the two *GbX* paralogs and the two α -globin paralogs have not been previously documented in any vertebrate taxon.

Phylogenetic Relationships

In broad outline, our reconstruction of phylogenetic relationships among vertebrate globins was consistent with previous studies (Hoffmann, Opazo, et al. 2010; Blank and Burmester 2012; Hoffmann, Opazo, Hoogewijs, et al. 2012; Hoffmann, Opazo, Storz, et al. 2012; Schwarze and Burmester 2013; Schwarze et al. 2014). Vertebrate globins were arranged into three separate clades, two representing the ancient *GbX* and *Ngb* gene lineages and one representing the

vertebrate-specific globins, including gnathostome α - and β -globin, *GbE*, *GbY*, and *Mb*, cyclostome *Hbs* and *Mbs*, and the *Cygb* genes of gnathostomes and cyclostomes (fig. 1). Support for monophyly of the three clades of vertebrate globins, expressed as Bayesian posterior probabilities, was higher than 0.99, and support for monophyly of the different vertebrate globin types was also high, with posterior probabilities higher than 0.99 in all cases except gnathostome α -globins (fig. 1). Compared with a recent phylogenetic analysis of vertebrate globins presented in Schwarze et al. (2014), the tree shown in figure 1 shows relatively strong support for the node joining cyclostome and gnathostome *Cygb*s (posterior probability = 0.91). Vertebrate-specific globins are arranged into three strongly supported clades containing 1) cyclostome *Hbs*, 2) cyclostome and gnathostome *Cygb*s, and 3) gnathostome α - and β -globin, *GbE*, *GbY*, and *Mb*.

Two of the nine elephant shark globins were placed in distinct *GbX* clades, and the remainder fell in well-supported clades of vertebrate-specific globins (fig. 1). None of the elephant shark globins was placed in the *Ngb* clade. For the majority of genes, phylogenetic placement of the elephant shark globins provided a clear means of inferring orthology; we identified two α -globin paralogs, two *GbX* paralogs, and single copies of β -globin, *GbY*, and *Mb*. Within the clade of vertebrate-specific globins, the position of the elephant shark globins matched expectations of the organismal phylogeny: the α/β -*Hb* and *Mb* genes of the elephant shark were grouped with corresponding sequences from other cartilaginous fish, and elephant shark *GbY* was placed sister to *GbY* sequences from bony vertebrates. Assigning orthologous relationships of the *GbX* sequences was more complicated, as one of the two elephant shark *GbX* paralogs was placed sister to cyclostome *GbX* genes, and the second was placed sister to a group that included teleost *GbX*s plus the coelacanth *GbX2* paralog (fig. 1).

A comparison of the organismal phylogeny with the phylogeny of *GbX* genes in figure 1 suggests the presence of multiple *GbX* paralogs in the common ancestor of vertebrates. To better address this issue, we performed a second phylogenetic analysis restricted to *GbX* paralogs, with increased taxonomic sampling of *GbX* sequences relative to the tree in figure 1, and using a group of relatively closely related acorn worm globins as outgroup. In this second analysis, *GbX* paralogs fell into four main clades: 1) *GbX* sequences from western clawed frog, teleost fish, plus one of the two *GbX* paralogs present in the gar, coelacanth, and elephant shark genomes, 2) *GbX* sequences from squamates and testudines, plus the *GbX1* paralogs of gar and coelacanth, 3) a group that includes all *GbX* paralogs of cyclostomes, and 4) the second elephant shark globin paralog, labeled as *GbX2* (fig. 2). Relationships among these lineages are well supported, and a comparison of the gene tree with the organismal phylogeny strongly suggests that the common ancestor of vertebrates possessed at least four *GbX* paralogs.

Patterns of Conserved Synteny

Previous studies of globin phylogeny and conserved synteny have shown that the diversification of vertebrate-specific




Fig. 1. Bayesian phylogram depicting relationships among globin genes from representative deuterostome taxa. Numbers next to the nodes are Bayesian posterior probabilities. Terminal branches leading to elephant shark globins are shown in red. The tree was rooted by using plant globins as outgroup sequences (see Materials and Methods). The inset on top shows the organismal phylogeny for the representative deuterostome taxa included in the analysis.




Fig. 2. Bayesian phylogram depicting relationships among *GbX* genes from representative vertebrate taxa. Numbers next to the nodes are Bayesian posterior probabilities. Patterns of conserved synteny are depicted for *GbX* paralogs in each of three clades.

globins is largely attributable to two rounds of whole-genome duplication that occurred in the stem lineage of vertebrates (Hoffmann, Opazo, Storz, et al. 2012). The hypothesis that the vertebrate ancestor possessed a total of four *GbX* paralogs is consistent with the expected outcome of two successive rounds of whole-genome duplication (Meyer and Schartl 1999). If the four *GbX* paralogs represent quadruplicated products of the “2 R” genome duplications in the vertebrate ancestor, then this should be reflected in a 4-fold pattern of intragenomic synteny. Specifically, each of the four *GbX* paralogs should be embedded in unlinked chromosomal segments that share similar, interdigitated arrangements of paralogous genes (Storz et al. 2013). To test this, we examined the physical linkage arrangements of the *GbX* paralogs and we characterized patterns of conserved synteny among vertebrates. We excluded teleost fish from this analysis because this group has undergone an additional round of whole-genome duplication, which greatly complicates synteny comparisons.

Our comparative genomic analyses revealed conserved synteny for two of the four *GbX* groups defined in the phylogeny (figs. 2 and 3). One of these *GbX* paralogs is flanked by *PLEKHG3* and *SRP14* in most species examined, whereas a

second one is flanked by *TIMM50* and *SUPT5H*. In coelacanth and spotted gar, this latter *GbX* paralog is also flanked by *PLEKHG2*, a paralog of *PLEKHG3*. Thus, the linked *PLEKHG* and *GbX* genes appear to have coduplicated, possibly during one or two rounds of whole-genome duplication prior to the divergence of cyclostomes and gnathostomes. With the notable exception of the *GbX1* sequences from lampreys, shared synteny was reflected in the phylogenetic reconstructions; the *GbX* paralogs that are flanked by *PLEKHG3* and *SRP14* on either side were phylogenetically placed with the *GbX* paralogs from teleost fish, whereas the *GbX* paralogs that are flanked by *TIMM50* and *SUPT5H* were placed with the *GbX* paralogs from squamates and testudines. The *GbX1* paralogs of lamprey deviate from this pattern, as they are flanked by *PLEKHG3* and *SRP14* on either side, but they were phylogenetically placed with another cyclostome paralog, *GbX2*. Although the inferred sister relationship between the cyclostome *GbX1* and *GbX2* paralogs has high statistical support (posterior probability = 0.99; fig. 2), such results should be interpreted cautiously due to the possible influence of base-composition bias (Qiu et al. 2011; Mehta et al. 2013; Smith et al. 2013) and other factors. Unfortunately, there is no synteny information for the cyclostome *GbX2*




Fig. 3. Patterns of conserved synteny in genomic regions that harbor paralogous *GbX* genes in representative vertebrate taxa.

genes or the elephant shark *GbX2* gene, as the cyclostome genes are found in isolated contigs and the elephant shark gene is not present in the genome assembly. Additional genomic data from cartilaginous fishes and cyclostomes will be required to assess which *GbX* paralogs might have originated via whole-genome duplication. Specifically, synteny information for the chromosomal region harboring orthologs of elephant shark *GbX2* in other cartilaginous fish may reveal whether this gene represents a product of the 2R whole-genome duplication that was secondarily lost in the ancestor of bony vertebrates.

Loss of *Ngb* in Chondrichthyes

All gnathostome vertebrates that have been investigated so far possess a *Ngb* gene, which is expressed in the nervous system. It was therefore surprising that *Ngb* was not detected in the genome or brain transcriptome of the elephant shark. The gene synteny *POMT2* - *Ngb* - *TMEM63C* - *ZDHHC22* is conserved among vertebrates; in the elephant shark, the genes *POMT2*, *TMEM63C*, and *ZDHHC22* were found on scaffold_319 in the expected order, but with *Ngb* missing. Further database searches did not reveal any evidence for a *Ngb* gene in the genome and transcriptome of the little skate (*Leucoraja erinacea*), or in the transcriptomes of the cloudy catshark (*Scyliorhinus torazame*), the spiny dogfish shark (*Squalus acanthias*), the nurse shark (*Ginglymostoma cirratum*), or the Pacific electric ray (*Torpedo californica*), suggesting that *Ngb* was lost the stem lineage of Chondrichthyes.

The Globin Gene Repertoire of the Vertebrate Common Ancestor

Comparative analysis of complete globin gene repertoires from representatives of all vertebrate classes indicates that

the last common ancestor of vertebrates possessed a repertoire of at least seven globin genes: single copies of *Adgb* and *Ngb*, four paralogous copies of *GbX*, and the single-copy progenitor of the entire set of vertebrate-specific globins (α/β -*Hb*, *Mb*, *GbE*, *GbY*, *Cygb*, and the cyclostome *Hbs* and *Mbs*). During the course of vertebrate evolution, *GbY*, *Mb*, *GbE*, *GbX*, and *Ngb* have each been lost in at least two different lineages (fig. 4). Given that *GbX* is present in the genomes of turtles and squamate reptiles (fig. 4), the principle of parsimony suggests that this gene must have been deleted independently in mammals and archosaurs (represented by birds and crocodilians). *Cygb* and *Adgb* are the only globins that have been retained in all vertebrate taxa examined to date.

Divergent Expression Domains of Paralogous Globins

Some of the elephant shark globins displayed expression profiles that were consistent with experimental results from orthologous globins in bony vertebrates. For example, *Adgb* was preferentially expressed in testis (fig. 5A), consistent with experimental results for mammals (Hoogewijs et al. 2012). *Cygb* was expressed in a broad range of tissues, including brain (fig. 5B), although overall expression levels were low relative to other bony vertebrates (Nakatani et al. 2004; Schmidt et al. 2004; Motoyama et al. 2014). *GbY* was expressed in a similarly broad range of tissues (fig. 5C), consistent with data from *Xenopus* (Fuchs et al. 2006). In contrast, the elephant shark *Mb*, α -globin, and *GbX* genes exhibited highly unexpected expression patterns.

Although *Mb* is most highly expressed in the myocytes of cardiac and skeletal muscle in most bony vertebrates, elephant shark *Mb* is expressed at a high level only in heart, but not in skeletal muscle (fig. 5D). The *Mb* gene has been deleted in anuran amphibians (Fuchs et al. 2006; Hoffmann et al. 2011), and it has been hypothesized that *Cygb* performs a compensatory *Mb*-like O_2 -storage function in skeletal muscle (Xi et al. 2007). In elephant shark there is no detectable expression of *Cygb* in skeletal muscle (fig. 5B), so the compensatory mechanism hypothesized for anuran amphibians is clearly not operative in elephant shark. It may be that the ancestral function of *Mb* was to facilitate oxygen delivery to cardiac myocytes in early gnathostomes that had single-circuit circulatory systems. The prominent expression of *Mb* in the skeletal muscle of many gnathostomes may have been a later innovation that coincided with the evolution of higher rates of aerobic metabolism and the concomitant increase in tissue oxygen demands.

One of the elephant shark α -globin paralogs ($\alpha 1$) and the sole β -globin gene showed highest expression in heart and spleen (fig. 5E and G), as expected for subunit isoforms of a tetrameric $\alpha_2\beta_2$ red blood cell Hb. Surprisingly, the other α -globin paralog ($\alpha 2$) was most highly expressed in the brain (fig. 5F). Expression of heterotetrameric Hb in brain neurons has been reported in mammals (Humphries et al. 1976; Ohyagi et al. 1994; Russo et al. 2013) and is associated with an enhanced neuronal oxygen supply (Schelshorn et al. 2009; Biagioli et al. 2009). The expression of a single Hb chain in




FIG. 4. Phyletic distribution of paralogous globin genes in the genomes of vertebrates.




FIG. 5. Expression profiles of elephant shark globin genes in ten tissues. Gene-specific mRNA levels were quantified using RNA-seq. Transcript abundances are measured in transcripts per million (TPM).

the shark brain provides evidence that this globin acts as a monomer, which excludes an Hb-like physiological function involving cooperative oxygen binding. It is intriguing that the brain-specific expression of this Hb chain is associated with deletion of the *Ngb* gene, and it suggests the hypothesis that α -globin 2 was co-opted to perform the brain-specific physiological functions that are performed by *Ngb* in bony vertebrates.

The two elephant shark *GbX* paralogs exhibited a surprising divergence in expression domains. Although *GbX1* was expressed in a diverse range of tissues, *GbX2* was most highly expressed in ovary and testis (fig. 5H and I). Studies of zebrafish and *Xenopus* documented predominant expression of *GbX* in the nervous system (Fuchs et al. 2006; Blank, Wollberg et al. 2011). However, those studies did not examine gene expression in the gonads, which suggests that the main

site of *GbX* expression may have been overlooked. Alternatively, *GbX* genes may have evolved different expression domains in cartilaginous fish and bony vertebrates.

Evolution of Vertebrate Globins

Historically, much of the uncertainty about the phylogenetic relationships among vertebrate globins was attributable to inadequate sampling of taxa and/or paralogous gene lineages. In recent years, the inclusion of a broader diversity of vertebrate globins in phylogenetic analyses has led to surprising findings regarding the antiquity of “taxon-specific” globins like *GbE*, which often turn out to be older than originally suspected (Hoffmann et al. 2011; Schwarze and Burmester 2013), and the independent origins of functionally analogous globins in disparate taxa that were previously assumed to be orthologous. For example, it was previously assumed that *Mb* and the single-copy progenitor of the α/β -Hbs originated via duplication of an ancestral, single-copy globin gene prior to the divergence between cyclostomes and gnathostomes, such that the ancestors of these two groups inherited orthologous copies of the same “proto-*Hb*” gene (Goodman et al. 1975, 1987). According to this scenario, the *Hb* genes of cyclostomes would be sister to the clade of gnathostome α - and β -globin genes: (*Mb* [cyclostome *Hb*, gnathostome *Hb*]). Only by including the full panoply of cyclostome-specific globins in phylogenetic analyses was it possible to infer that functionally analogous “Hbs” and “Mbs” evolved independently from different precursor globins that were present in the vertebrate ancestor (Hoffmann, Opazo, et al. 2010; Schwarze et al. 2014). Similarly, because of extensive lineage-specific gene losses, the single copy *GbX* genes of frog, most teleost fish, testudines, and squamates were thought to be 1:1 orthologs, and the possession of two copies in coelacanth was attributed to lineage-specific duplication (Schwarze and Burmester 2013). However, the inclusion of globin sequences from elephant shark, coelacanth, and lampreys in our analyses suggests a more complex historical scenario. The phylogenetic reconstructions and patterns of conserved synteny suggest that whole-genome duplication and/or smaller scale duplication events produced a set of four unlinked *GbX* paralogs in the common ancestor of vertebrates. During the ensuing approximately 500 My of vertebrate evolution, lineage-specific gene losses led to a gradual winnowing of *GbX* diversity, with different lineages retaining different *GbX* paralogs.

Materials and Methods

DNA Sequence Data

We characterized the complete globin gene repertoire of the elephant shark (*C. milii*) by assembling RNA-seq reads from ten tissues (brain, gills, heart, intestine, kidney, liver, muscle, ovary, spleen, and testis; Venkatesh et al. 2014). We then used the identified transcripts to search against databases of known vertebrate globins. We also used the elephant shark transcripts to search against the nonredundant (NR) protein database available at NCBI using BLASTX to confirm their identities as authentic globins. We then mapped all confirmed globin transcripts to the elephant shark genome

assembly (Assembly Callorhinchus_milii-6.1.3, available at <http://esharkgenome.imcb.a-star.edu.sg/>) using BLASTN, where full-length coding sequences were predicted by a combination of similarity of the transcripts to the genome sequence (BLASTN) and similarity of the genome sequence to known globin protein sequences from the NCBI database (BLASTX). The predicted protein sequences were further refined by manual inspection of exon–intron boundaries. One of the transcripts, which is full-length and encodes *GbX2*, had no match in the genome assembly. This gene is presumably present in the gaps of the genome assembly. Our searches in the NCBI database identified an additional globin gene, α -globin 2, which was previously identified in the full-length cDNA data set (accession number AFK11062; Tan et al. 2012) but is not present in the current genome assembly.

Our comprehensive analyses also included the complete globin gene repertoire of 19 deuterostome taxa, including 17 chordates (14 vertebrates, two urochordates, and one cephalochordate), one hemichordate and one echinoderm. The vertebrate species included three mammals (human, *Homo sapiens*; gray short-tailed opossum, *Monodelphis domestica*; and platypus, *O. anatinus*), three birds (chicken, *Gallus gallus*; turkey, *Meleagris gallopavo*; and zebra finch, *Taeniopygia guttata*), one squamate reptile (green anole lizard, *A. carolinensis*), one amphibian (western clawed frog, *Xenopus tropicalis*), one lobe-finned fish (West Indian Ocean coelacanth, *L. chalumnae*), and five teleost fish (fugu, *Takifugu rubripes*; medaka, *Oryzias latipes*; green-spotted pufferfish, *Tetraodon nigroviridis*; three-spined stickleback, *Gasterosteus aculeatus*; and zebrafish, *Danio rerio*). The nonvertebrate species included two sea squirts (*Ciona intestinalis* and *C. savignyi*, Urochordata), amphioxus (*Branchiostoma floridae*, Cephalochordata), the acorn worm (*Saccoglossus kowalevskii*, Hemichordata), and the purple sea urchin (*Strongylocentrotus purpuratus*, Echinodermata). Because the sea urchin possesses a globin gene that encodes 16 putative globin domains (Bailey and Vinogradov 2008), we considered each of the domains as separately alignable globin sequences in our analyses. In addition to retrieving the complete globin gene repertoires of the species listed above, we also included individual globin sequences from several additional taxa. For echinoderms, we included sequences from two subunits of the coelomic red blood cell Hbs of the sea cucumber, *Caudina arenicola*. For cartilaginous fish, we obtained α - and β -*Hb* sequences from the red stingray (*Dasyatis akajei*) and gummy houndshark (*Mustelus antarcticus*), as well as *Mb* sequences from the gummy houndshark and the Port Jackson shark (*Heterodontus portusjacksoni*). For birds, we included the *GbE* sequence of the mallard duck (*Anas platyrhynchos*). For cyclostomes, we included the *GbX1*, *GbX2*, and *Cygb* genes of the Arctic and sea lampreys (*Lethenteron japonicum* and *Petromyzon marinus*, respectively), the *Cygb* of the European brook lamprey (*Lampetra planeri*), and the *Hbs* of the arctic lamprey and Atlantic hagfish (*Myxine glutinosa*; Schwarze et al. 2014). In order to include representatives from each of the main deuterostome lineages, we also included two globin sequences from the acoel *Symsagittifera roscoffensis*,

three globin sequences from the nemertodermatid *Nemertoderma westbladi*, plus one sequence from the xenoturbellid, *Xenoturbella bocki*. Nine plant globin sequences were included as outgroup. The full list of sequences and the corresponding accession numbers are presented in [supplementary table S1, Supplementary Material](#) online. The order of the helical domains of the globin domain of *Androglobin* has been rearranged relative to all other globins in this study (Hoogewijs et al. 2012). Because of the corresponding alignment challenges, this sequence was not included in the phylogenetic analyses.

For the analyses that focused on resolving orthologous relationships among vertebrate *GbX* paralogs, we also included additional *GbX* genes from other vertebrate groups to increase our taxonomic coverage. We added *GbX* paralogs from an holostean fish (spotted gar, *Lepisosteus oculatus*), three testudines (green sea turtle, *Chelonia mydas*; painted turtle, *Chrysemys picta*; and Chinese softshell turtle, *Pelodiscus sinensis*), two squamates (Burmese python, *Python morulus*; and anole lizard, *A. carolinensis*), and additional teleost fish (Atlantic cod, *Gadus morhua*; tilapia, *Oreochromis niloticus*; goldfish, *Carassius auratus*; platyfish, *Xiphophorus maculatus*; and cave fish, *Astyanax mexicanus*). For the sake of providing comprehensive coverage of amniotes, we also surveyed genome assemblies and sequence databases to identify globin genes in crocodylians (Green et al. 2014) and birds (Grispo et al. 2012; Zhang et al. 2014; Opazo et al. 2015) using the bioinformatic strategies outlined above.

Phylogenetic Analyses

We estimated phylogenetic relationships among globin genes of deuterostomes using a Bayesian approach, as implemented in the program MrBayes v3.1.2 (Ronquist and Huelsenbeck 2003) under a mixed model of amino acid substitution. Sequence alignment was carried out using the G-INS-i strategy from MAFFT v.6 (Katoh et al. 2009). Two simultaneous independent runs were performed for 10×10^6 iterations of a Markov Chain Monte Carlo algorithm, with six simultaneous chains sampling trees every 1,000 generations. Support for the nodes and parameter estimates were derived from a majority rule consensus of the last 5,000 trees sampled after convergence. The average standard deviation of split frequencies remained 0.01 after the burn-in threshold.

Gene Expression Analyses

Paired-end RNA-seq data for ten adult elephant shark tissues (brain, gills, heart, intestine, kidney, liver, muscle, ovary, spleen, and testis) were obtained from SRA accession SRA054255 (Venkatesh et al. 2014). Sequence reads were aligned against the elephant shark transcriptome (NR set of ~18,900 transcripts; Venkatesh et al. 2014) using Bowtie v. 0.12.9 (Langmead et al. 2009). Transcript abundances were estimated using RSEM v1.2.3 (Li and Dewey 2011). Default settings were used for both programs. Transcript abundances are measured in transcripts per million.

Supplementary Material

Supplementary table S1 is available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

Acknowledgments

This work was funded by grants from the National Institutes of Health/National Heart, Lung, and Blood Institute to J.F.S. (R01 HL087216 and HL087216-S1), grants from the National Science Foundation to J.F.S. (IOS-0949931), and F.G.H. (EPS-0903787), a grant from Fondo Nacional de Desarrollo Científico y Tecnológico to J.C.O. (FONDECYT 1120032), a grant by the Deutsche Forschungsgemeinschaft to T.B. (BU 956/18-1), and a grant from the Biomedical Research Council of A*STAR to B.V. The authors thank the associate editor and two anonymous reviewers for helpful suggestions.

References

- Awenius C, Hankeln T, Burmester T. 2001. Neuroglobins from the zebrafish *Danio rerio* and the pufferfish *Tetraodon nigroviridis*. *Biochem Biophys Res Commun*. 287:418–421.
- Bailey X, Vinogradov S. 2008. The bilaterian sea urchin and the radial starlet sea anemone globins share strong homologies with vertebrate neuroglobins. In: Bolognesi M, di Prisco G, Verde C, editors. *Dioxygen binding and sensing proteins*. New York: Springer Verlag, p. 191–201.
- Biagioli M, Pinto M, Cesselli D, Zaninello M, Lazarevic D, Roncaglia P, Simone R, Vlachouli C, Plessy C, Bertin N, et al. 2009. Unexpected expression of alpha- and beta-globin in mesencephalic dopaminergic neurons and glial cells. *Proc Natl Acad Sci U S A*. 106: 15454–15459.
- Blank M, Burmester T. 2012. Widespread occurrence of N-terminal acylation in animal globins and possible origin of respiratory globins from a membrane-bound ancestor. *Mol Biol Evol*. 29:3553–3561.
- Blank M, Kiger L, Thielebein A, Gerlach F, Hankeln T, Marden MC, Burmester T. 2011. Oxygen supply from the bird's eye perspective: globin E is a respiratory protein in the chicken retina. *J Biol Chem*. 286:26507–26515.
- Blank M, Wollberg J, Gerlach F, Reimann K, Roesner A, Hankeln T, Fago A, Weber RE, Burmester T. 2011. A membrane-bound vertebrate globin. *PLoS One* 6:e25292.
- Burmester T, Ebner B, Weich B, Hankeln T. 2002. Cytochrome b: a novel globin type ubiquitously expressed in vertebrate tissues. *Mol Biol Evol*. 19:416–421.
- Burmester T, Haberkamp M, Mitz SA, Roesner A, Schmidt M, Ebner B, Gerlach F, Fuchs C, Hankeln T. 2004. Neuroglobin and cytochrome b: genes, proteins and evolution. *IUBMB Life* 56:703–707.
- Burmester T, Hankeln T. 2009. What is the function of neuroglobin? *J Exp Biol*. 212 1423–1428.
- Burmester T, Hankeln T. 2014. Function and evolution of vertebrate globins. *Acta Physiol*. 211:501–514.
- Burmester T, Weich B, Reinhardt S, Hankeln T. 2000. A vertebrate globin expressed in the brain. *Nature* 407:520–523.
- Cossins AR, Williams DR, Foulkes NS, Berenbrink M, Kipar A. 2009. Diverse cell-specific expression of myoglobin isoforms in brain, kidney, gill and liver of the hypoxia-tolerant carp and zebrafish. *J Exp Biol*. 212:627–638.
- Dröge J, Makalowski W. 2011. Phylogenetic analysis reveals wide distribution of globin X. *Biol Direct*. 6:54.
- Fuchs C, Burmester T, Hankeln T. 2006. The amphibian globin gene repertoire as revealed by the *Xenopus* genome. *Cytogenet Genome Res*. 112:296–306.
- Fuchs C, Heib V, Kiger L, Haberkamp M, Roesner A, Schmidt M, Hamdane D, Marden MC, Hankeln T, Burmester T. 2004. Zebrafish reveals different and conserved features of vertebrate

- neuroglobin gene structure, expression pattern, and ligand binding. *J Biol Chem.* 279:24116–24122.
- Fuchs C, Luckhardt A, Gerlach F, Burmester T, Hankeln T. 2005. Duplicated cytoglobin genes in teleost fishes. *Biochem Biophys Res Commun.* 337:216–223.
- Goodman M, Czelusniak J, Koop BF, Tagle DA, Slightom JL. 1987. Globins: a case study in molecular phylogeny. *Cold Spring Harb Symp Quant Biol.* 52:875–890.
- Goodman M, Moore GW, Matsuda G. 1975. Darwinian evolution in the genealogy of haemoglobin. *Nature* 253:603–608.
- Green RE, Braun EL, Armstrong J, Earl D, Nguyen N, Hickey G, Vandeweghe MW, St John JA, Capella-Gutiérrez S, Castoe TA, et al. 2014. Three crocodylian genomes reveal ancestral patterns of evolution among archosaurs. *Science* 346:1254449.
- Grispo MT, Natarajan C, Projecto-García J, Moriyama H, Weber RE, Storz JF. 2012. Gene duplication and the evolution of hemoglobin isoform differentiation in birds. *J Biol Chem.* 287:37647–37658.
- Hankeln T, Burmester T. 2008. Neuroglobin and cytoglobin. In: Gosh A, editor. The smallest biomolecules: diatomics and their interactions with heme proteins. Amsterdam: Elsevier. p. 203–218.
- Hankeln T, Ebner B, Fuchs C, Gerlach F, Haberkamp M, Laufs TL, Roesner A, Schmidt M, Weich B, Wystub S, et al. 2005. Neuroglobin and cytoglobin in search of their role in the vertebrate globin family. *J Inorg Biochem.* 99:110–119.
- Hedges SB. 2009. Vertebrates (Vertebrata). In: Hedges SB, Kumar S, editors. The timetree of life. New York: Oxford University Press. p. 309–314.
- Helbo S, Weber RE, Fago A. 2013. Expression patterns and adaptive functional diversity of vertebrate myoglobins. *Biochim Biophys Acta.* 1834:1832–1839.
- Hoffmann FG, Opazo JC, Hoogewijs D, Hankeln T, Ebner B, Vinogradov SN, Bailly X, Storz JF. 2012. Evolution of the globin gene family in deuterostomes: lineage-specific patterns of diversification and attrition. *Mol Biol Evol.* 29:1735–1745.
- Hoffmann FG, Opazo JC, Storz JF. 2010. Gene cooption and convergent evolution of oxygen transport hemoglobins in jawed and jawless vertebrates. *Proc Natl Acad Sci U S A.* 107:14274–14279.
- Hoffmann FG, Opazo JC, Storz JF. 2011. Differential loss and retention of cytoglobin, myoglobin, and globin-E during the radiation of vertebrates. *Genome Biol Evol.* 3:588–600.
- Hoffmann FG, Opazo JC, Storz JF. 2012. Whole-genome duplications spurred the functional diversification of the globin gene superfamily in vertebrates. *Mol Biol Evol.* 29:303–312.
- Hoffmann FG, Storz JF, Gorr TA, Opazo JC. 2010. Lineage-specific patterns of functional diversification in the α - and β -globin gene families of tetrapod vertebrates. *Mol Biol Evol.* 27:1126–1138.
- Hoogewijs D, Ebner B, Germani F, Hoffmann FG, Fabrizius A, Moens L, Burmester T, Dewilde S, Storz JF, Vinogradov SN. 2012. Androglobin: a chimeric globin in metazoans that is preferentially expressed in mammalian testes. *Mol Biol Evol.* 29:1105–1114.
- Humphries S, Windass J, Williamson R. 1976. Mouse globin gene expression in erythroid and non-erythroid tissues. *Cell* 7:267–277.
- Inoue JG, Miya M, Lam K, Tay B-H, Danks JA, Bell J, Walker TI, Venkatesh B. 2010. Evolutionary origin and phylogeny of the modern holocephalans (Chondrichthyes: Chimaeriformes): a mitogenome perspective. *Mol Biol Evol.* 27:2576–2586.
- Kakar S, Hoffmann FG, Storz JF, Fabian M, Hargrove MS. 2010. Structure and reactivity of hexacoordinate hemoglobins. *Biophys Chem.* 152:1–14.
- Katoh K, Asimenos G, Toh H. 2009. Multiple alignment of DNA sequences with MAFFT. *Methods Mol Biol.* 537:39–64.
- Kugelstadt D, Haberkamp M, Hankeln T, Burmester T. 2004. Neuroglobin, cytoglobin, and a novel, eye-specific globin from chicken. *Biochem Biophys Res Commun.* 325:719–725.
- Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.
- Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12:323.
- Mehta TK, Ravi V, Yamasaki S, Lee AP, Lian MM, Tay B, Tohari S, Yanai S, Tay A, Brenner S, et al. 2013. Evidence for at least six Hox clusters in the Japanese lamprey (*Lethenteron japonicum*). *Proc Natl Acad Sci U S A.* 110:16044–16049.
- Meyer A, Schartl M. 1999. Gene and genome duplications in vertebrates: the one-to-four (-to-eight in fish) rule and the evolution of novel gene functions. *Curr Opin Cell Biol.* 11:699–704.
- Motoyama H, Komiya T, Thuy le TT, Tamori A, Enomoto M, Morikawa H, Iwai S, Uchida-Kobayashi S, Fujii H, Hagihara A, et al. 2014. Cytoglobin is expressed in hepatic stellate cells, but not in myofibroblasts, in normal and fibrotic human liver. *Lab Invest.* 94:192–207.
- Nakatani K, Okuyama H, Shimahara Y, Saeki S, Kim DH, Nakajima Y, Seki S, Kawada N, Yoshizato K. 2004. Cytoglobin/STAP, its unique localization in splanchnic fibroblast-like cells and function in organ fibrogenesis. *Lab Invest.* 84:91–101.
- Nishi H, Inagi R, Kato H, Tanemoto M, Kojima I, Son D, Fujita T, Nangaku M. 2008. Hemoglobin is expressed by mesangial cells and reduces oxidant stress. *J Am Soc Nephrol.* 19:1500–1508.
- Ohyagi Y, Yamada T, Goto I. 1994. Hemoglobin as a novel protein developmentally regulated in neurons. *Brain Res.* 635:323–327.
- Opazo JC, Butts GT, Nery MF, Storz JF, Hoffmann FG. 2013. Whole-genome duplication and the functional diversification of teleost fish hemoglobins. *Mol Biol Evol.* 30:140–153.
- Opazo JC, Hoffmann FG, Witt CC, Berenbrink M, Storz JF. Forthcoming 2015. Gene turnover in the avian globin gene families and evolutionary changes in hemoglobin isoform expression. *Mol Biol Evol.* doi: 10.1093/molbev/msu341.
- Pesce A, Bolognesi M, Bocedi A, Ascenzi P, Dewilde S, Moens L, Hankeln T, Burmester T. 2002. Neuroglobin and cytoglobin. Fresh blood for the vertebrate globin family. *EMBO Rep.* 3:1146–1151.
- Qiu H, Hildebrand F, Kuraku S, Meyer A. 2011. Unresolved orthology and peculiar coding sequence properties of lamprey genes: the KCNA gene family as test case. *BMC Evol Biol.* 12:325.
- Qiu Y, Sutton L, Riggs AF. 1998. Identification of myoglobin in human smooth muscle. *J Biol Chem.* 273:23426–23432.
- Roesner A, Fuchs C, Hankeln T, Burmester T. 2005. A globin gene of ancient evolutionary origin in lower vertebrates: evidence for two distinct globin families in animals. *Mol Biol Evol.* 22:12–20.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
- Russo R, Zucchelli S, Codrich M, Marcuzzi F, Verde C, Gustincich S. 2013. Hemoglobin is present as a canonical $\alpha_2\beta_2$ tetramer in dopaminergic neurons. *Biochim Biophys Acta.* 1834:1939–1943.
- Schelshorn DW, Schneider A, Kuschinsky W, Weber D, Kruger C, Dittgen T, Burgers HF, Sabouri F, Gassler N, Bach A, et al. 2009. Expression of hemoglobin in rodent neurons. *J Cereb Blood Flow Metab.* 29:585–595.
- Schmidt M, Gerlach F, Avivi A, Laufs T, Wystub S, Simpson JC, Nevo E, Saaler-Reinhardt S, Reuss S, Hankeln T, et al. 2004. Cytoglobin is a respiratory protein in connective tissue and neurons, which is up-regulated by hypoxia. *J Biol Chem.* 279:8063–8069.
- Schwarze K, Burmester T. 2013. Conservation of globin genes in the “living fossil” *Latimeria chalumnae* and reconstruction of the evolution of the vertebrate globin family. *Biochim Biophys Acta.* 1834:1801–1812.
- Schwarze K, Campbell KL, Hankeln T, Storz JF, Hoffman FG, Burmester T. 2014. The globin gene repertoire of lampreys: convergent evolution of hemoglobin and myoglobin in jawed and jawless vertebrates. *Mol Biol Evol.* 31:2708–2721.
- Smith JS, Kuraku S, Holt C, Sanka-Spengler T, Jiang N, Campbell MS, Yandell MD, Manousaki T, Meyer A, Bloom OE, et al. 2013. Sequencing of the sea lamprey (*Petromyzon marinus*) genome provides insights into vertebrate evolution. *Nat Genet.* 45:415–421.

- Storz JF, Opazo JC, Hoffmann FG. 2011. Phylogenetic diversification of the globin gene superfamily in chordates. *IUBMB Life* 63:313–322.
- Storz JF, Opazo JC, Hoffmann FG. 2013. Gene duplication, genome duplication, and the functional diversification of vertebrate globins. *Mol Phylogenet Evol.* 66:469–478.
- Straub AC, Lohman AW, Billaud M, Johnstone SR, Dwyer ST, Lee MY, Bortz PS, Best AK, Columbus L, Gaston B, et al. 2012. Endothelial cell expression of haemoglobin alpha regulates nitric oxide signalling. *Nature* 491:473–477.
- Tan YY, Kodzius R, Tay B-H, Tay A, Brenner S, Venkatesh B. 2012. Sequencing and analysis of full-length cDNAs, 5'-ESTs and 3'-ESTs from a cartilaginous fish, the elephant shark (*Callorhynchus milii*). *PLoS One* 7:e47174.
- Trent JT, Hargrove MS. 2002. A ubiquitously expressed human hexacoordinate hemoglobin. *J Biol Chem.* 277:19538–19545.
- Venkatesh B, Kirkness EF, Loh YH, Halpern AL, Lee AP, Johnson J, Dandona N, Viswanathan, Tay A, Venter JC, et al. 2007. Survey sequencing and comparative analysis of the elephant shark (*Callorhynchus milii*) genome. *PLoS Biol.* 5:e101.
- Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, Ohta Y, Flajnik MF, Sutoh Y, Kasahara M, et al. 2014. Elephant shark genome provides unique insights into gnathostome evolution. *Nature* 505: 174–179.
- Wittenberg JB, Wittenberg BA. 2003. Myoglobin function reassessed. *J Exp Biol.* 206:2011–2020.
- Wystub S, Ebner B, Fuchs C, Weich B, Burmester T, Hankeln T. 2004. Interspecies comparison of neuroglobin, cytoglobin and myoglobin: sequence evolution and candidate regulatory elements. *Cytogenet Genome Res.* 105:65–78.
- Xi Y, Obara M, Ishida Y, Ikeda S, Yoshizato K. 2007. Gene expression and tissue distribution of cytoglobin and myoglobin in the Amphibia and Reptilia: possible compensation of myoglobin with cytoglobin in skeletal muscle cells of anurans that lack the myoglobin gene. *Gene* 398:94–102.
- Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, Storz JF, Antunes A, Greenwold MJ, Meredith RW, et al. 2014. Comparative genomics reveals insights into avian genome evolution and adaptation. *Science* 346:1311–1320.